



**QUEEN'S
UNIVERSITY
BELFAST**

A Coordination and Ligand Replacement Based Three-Input Colorimetric Logic Gate Sensing Platform for Melamine, Mercury Ions, and Cysteine

Zhang, L., Yuan, Y., Wen, X., Li, Y., Cao, C., & Xiong, Q. (2015). A Coordination and Ligand Replacement Based Three-Input Colorimetric Logic Gate Sensing Platform for Melamine, Mercury Ions, and Cysteine. *RSC Advances*, 5(73), 59106-59133. <https://doi.org/10.1039/C5RA09570K>

Published in:
RSC Advances

Document Version:
Peer reviewed version


Queen's University Belfast - Research Portal:
[Link to publication record in Queen's University Belfast Research Portal](#)

Publisher rights
© Royal Society of Chemistry 2015

General rights
Copyright for the publications made accessible via the Queen's University Belfast Research Portal is retained by the author(s) and / or other copyright owners and it is a condition of accessing these publications that users recognise and abide by the legal requirements associated with these rights.

Take down policy
The Research Portal is Queen's institutional repository that provides access to Queen's research output. Every effort has been made to ensure that content in the Research Portal does not infringe any person's rights, or applicable UK laws. If you discover content in the Research Portal that you believe breaches copyright or violates any law, please contact openaccess@qub.ac.uk.

RSC Advances



This article can be cited before page numbers have been issued, to do this please use: L. Zhang, Y. Yuan, X. Wen, Y. Li, C. Cao and Q. Xiong, *RSC Adv.*, 2015, DOI: 10.1039/C5RA09570K.



This is an *Accepted Manuscript*, which has been through the Royal Society of Chemistry peer review process and has been accepted for publication.

Accepted Manuscripts are published online shortly after acceptance, before technical editing, formatting and proof reading. Using this free service, authors can make their results available to the community, in citable form, before we publish the edited article. This *Accepted Manuscript* will be replaced by the edited, formatted and paginated article as soon as this is available.

You can find more information about *Accepted Manuscripts* in the [Information for Authors](#).

Please note that technical editing may introduce minor changes to the text and/or graphics, which may alter content. The journal's standard [Terms & Conditions](#) and the [Ethical guidelines](#) still apply. In no event shall the Royal Society of Chemistry be held responsible for any errors or omissions in this *Accepted Manuscript* or any consequences arising from the use of any information it contains.



Journal Name

ARTICLE

A Coordination and Ligand Replacement Based Three-Input Colorimetric Logic Gate Sensing Platform for Melamine, Mercury Ions, and Cysteine

Lulu Zhang^a, Yanwen Yuan^a, Xinglin Wen^a, Yue Li^b, Cuong Cao^{c*}, and Qihua Xiong^{a,d*}Received 00th January 20xx,
Accepted 00th January 20xx

DOI: 10.1039/x0xx00000x

www.rsc.org/

Discrimination of different species in various target scopes within a single sensing platform can provide many advantages such as simplicity, rapidness, and cost effectiveness. Here we design a three-input colorimetric logic gate based on the aggregation and anti-aggregation of gold nanoparticles (Au NPs) for the sensing of melamine, cysteine, and Hg²⁺. The concept takes advantages of the highly specific coordination and ligand replacement reactions between melamine, cysteine, Hg²⁺, and Au NPs. Different outputs are obtained with the combinational inputs in the logic gates, which can serve as a reference to discriminate different analytes within a single sensing platform. Furthermore, besides the intrinsic sensitivity and selectivity of Au NPs to melamine-like compounds, the "INH" gates of melamine/cysteine and melamine/Hg²⁺ in this logic system can be employed for sensitive and selective detections of cysteine and Hg²⁺, respectively.

Introduction

Maintaining a healthy life has become one of the most pressing concerns across the world for decades, especially with the increasing issues on food safety, environmental contamination, and personal healthcare. In particular, the shocking scandals of melamine in infant formula in 2008, and in yoghurt candy in 2014 have further warned the general public that food safety is a long-lasting battle for the human society.^{1, 2} Heavy metal contaminations, such as mercury, on the other hand, represents one of the most severe environmental challenges.³ As a sulphur-containing amino acid, cysteine plays a crucial biological role in the human body. It is also a potential neurotoxin at elevated levels,^{4, 5} a biomarker for various medical conditions,^{6, 7} and a disease-associated physiological regulator.⁸⁻¹⁰ Therefore, a rapid and low-cost method for identification and detection of these targets is of crucial importance for environmental protection, food safety and medical diagnostics as well as for fundamental researches. However, conventional methods employed in the sensing of Hg²⁺ (e.g. cold-vapor atomic fluorescence spectrometry,¹¹ cold-vapor atomic absorption spectrometry,¹² inductively coupled

plasma-mass spectrometry,¹³ and X-ray absorption spectroscopy¹⁴), melamine (e.g. high-performance liquid chromatography,¹⁵ and mass spectroscopy¹⁶), and cysteine (e.g. electrochemical voltammetry,^{17, 18} and fluorescence-coupled HPLC techniques^{19, 20}) are often expensive, labour-intensive, time-consuming, and non-portable. Therefore, low cost, simple, rapid, and portable methods are highly desired.

In this context, colorimetric sensing based on analytes induced aggregation or anti-aggregation of Au NPs has been widely used in chemo/biosensors for the detection of various analytes owing to its integration of simplicity, rapidness and cost effectiveness.²¹⁻²⁷ In addition, the aggregation or anti-aggregation processes of Au NPs are often accompanied by a shift of the surface plasmon band as well as a clear colour change between blue and red due to the inter-particle plasmon coupling,^{28, 29} which can be easily observed either by naked eye or simple spectroscopy measurements. However, every sensing method has its own strengths and weaknesses due to its own limitations. Up to now, most of the colorimetric platforms based on Au NPs are designed to detect one certain target and cannot be employed for the discrimination of different species in various target scopes within a single sensing platform.

Molecular logic gates which borrow the concept of logic gate to construct chemo/biosensors provide a possible way of solving the above problem. Depending on the number and the relationship of the inputs in the system, a wide range of logic functions which include YES, NOT, AND, OR, NOR, NAND, INH, and XOR could be obtained.³⁰ Analytes can be distinguished through different signal outputs upon changing the inputs. Therefore, different types of molecular sensors based on logic gate operations have been developed for the sensing of various targets such as metal ions,^{31, 32} neurotransmitter release,³³ DNA and aptamer,³⁴ and iodide.³⁵

^a Division of Physics and Applied Physics, School of Physical and Mathematical Sciences, Nanyang Technological University, Singapore 637371. E-mail: Qihua@ntu.edu.sg

^b Key Laboratory of Materials Physics, Institute of Solid State Physics, Chinese Academy of Sciences, Hefei, 230031, P.R. China

^c Institute for Global Food Security, School of Biological Sciences, Queen's University Belfast, 18-30 Malone Road, Belfast, BT9 5BN, United Kingdom. E-mail: c.cao@qub.ac.uk

^d NOVITAS, Nanoelectronics Center of Excellence, School of Electrical and Electronic Engineering, Nanyang Technological University, Singapore 639798

† Electronic Supplementary Information (ESI) available. See DOI: 10.1039/x0xx00000x

ARTICLE

Journal Name

To date, colorimetric logic gates based on the aggregation or anti-aggregation of Au NPs are mostly formulated by two inputs due to the lack of highly specific interactions between multiple target species,^{24, 36-40} which thereby hampers the construction of logically inter-connected sensing system with more than two inputs. Furthermore, the input species lack highly specific interactions with each other or Au NPs may also lead to the uncontrollable behaviour of Au NPs. Therefore, ensuring highly specific interactions between the introduced species which serve as inputs plays a key role in rationally designing a logic gate system with more inputs. Herein, we report a new three-input molecular logic gate for the sensing of melamine, cysteine, and Hg^{2+} based on the mechanism of analyte-induced aggregation and anti-aggregation of Au NPs. The aggregation and anti-aggregation of Au NPs can be well controlled due to the highly specific coordination and ligand replacement reactions between melamine, cysteine, Hg^{2+} , and Au NPs. Therefore, a variety of inter-connected logic gates with different logic operations which include AND, OR, and INH can be built depending on the type of interactions between the targeted species and Au NPs. The highly specific interactions between the targeted species and Au NPs endow the as designed system with high selectivity to the targets. Furthermore, the intrinsic sensitivity of Au NPs to melamine, which has been carefully studied before,^{23, 41} enables the inter-connected sensing system to have good sensitivity to cysteine, and Hg^{2+} .

Experimental section

Materials and measurements

$\text{Hg}(\text{ClO}_4)_2 \cdot \text{H}_2\text{O}$, $\text{CrCl}_3 \cdot 6\text{H}_2\text{O}$, KCl, cyromazine, cyanuric acid, dicyandiamide, butylamine, acetonitrile, glycine, alanine, serine, proline, aspartic acid, asparagine, glutamic acid, ornithine, glutamine, histidine, tyrosine and arginine were of analytical grade and obtained from Sigma-Aldrich, Singapore. Cysteine, melamine, sodium citrate anhydrous, $\text{HAuCl}_4 \cdot 3\text{H}_2\text{O}$, ZnCl_2 , PbCl_2 , $\text{CuCl}_2 \cdot 2\text{H}_2\text{O}$, CdCl_2 , FeCl_3 , and NiCl_2 were purchased from Alfa Aesar. Ultrapure water (18 M Ω) was used for all solution preparations. UV-vis absorption spectra were acquired by using a PerkinElmer LAMBDA 950 spectrophotometer. TEM was performed on a JEOL1400 transmission electron microscope with an accelerating voltage of 100 kV. SEM image was acquired on a scanning electron microscopy (SEM; JEOL JSM-7001F).

Synthesis of ~13 nm and ~30 nm Au NPs

~13 nm colloidal Au NPs were prepared by the reduction of HAuCl_4 with trisodium citrate in aqueous solution.²³ Typically, 10 mL of trisodium citrate (38.8 mM) was rapidly injected into a boiling solution of HAuCl_4 (100 mL, 1 mM), and the mixed solution was further refluxed for another 15 min into a wine-red suspension. The suspension was gradually cooled to room temperature under stirring. The concentration of the Au NPs as determined by UV-vis spectrometry was 10 nM. Then 10 mL of Au NPs suspension was diluted with 50 mL of deionised water to give a total volume of 60 mL as a stock liquid for the sensing

of analytes. ~30 nm colloidal Au NPs were prepared by a similar method. Typically, 0.25 mL of 0.1 M HAuCl_4 was added to 100 mL of ultrapure water and then heated to boiling under magnetic stirring. After quickly injecting 1.5 mL of 1% trisodium citrate, the mixed solution was refluxed for 30 min and the Au NPs were synthesized. Then 10 mL of the Au NPs solution was diluted with 10 mL of deionised water to give a total volume of 20 mL as a stock liquid for the sensing of analyte.

Operations of the colorimetric logic gates

For the logic gate operations, different analytes were mixed thoroughly first and then added into 1 mL of the above ~13 nm Au NPs suspension with the final effective concentration of the analytes at 6×10^{-5} M (melamine), 3.6×10^{-4} M (cysteine), and 1.8×10^{-4} M (Hg^{2+}). To eliminate the effect of dilution, the volume of the mixture of the analytes was fixed less than 30 μL . After 2 minutes, absorbance spectra of Au NPs solutions were recorded with UV-vis spectrometer. The photos of the above Au NPs solutions at the same time were also taken by a digital camera.

Sensing of melamine

For analysis, 10 μL of melamine aqueous solution with certain concentration was added into 1 mL of Au NPs solution to study the aggregation degree of Au NPs. For selectivity experiments, 10 μL of one of the different nitrogen containing compounds in aqueous solution was added into 1 mL of Au NPs solution, the final effective concentrations of different nitrogen containing compounds in Au NPs solutions are all 1.5×10^{-6} M. After mixing thoroughly for 2 minutes, absorbance spectra and photos of Au NPs solutions at the same time were recorded with UV-vis spectrometer and the digital camera, respectively.

Sensing of cysteine

For selectivity experiments, 10 μL of one of the different amino acids and 10 μL of melamine in aqueous solution were mixed thoroughly first and then added into 1 mL of Au NPs solution, the final effective concentrations of the different amino acids and melamine were all 6×10^{-5} M. After mixing thoroughly for 2 minutes, absorbance spectra and photos of Au NPs solutions at the same time were recorded with UV-Vis spectrometer and the digital camera, respectively. To study the limit of detection of cysteine, aggregation degree of 1 mL of Au NPs solution induced by melamine at a final concentration of 1.4×10^{-6} M was used as a control sample. Then 10 μL of cysteine at different concentrations and 10 μL of melamine in aqueous solution were mixed thoroughly first and then added into 1 mL of Au NPs solutions, the final effective concentration of cysteine were ranging from 0 to 3×10^{-6} M, whereas the final concentration of melamine were all 1.4×10^{-6} M. After mixing thoroughly for 2 minutes, absorbance spectra and photos of Au NPs solutions at the same time were recorded with UV-vis spectrometer and the digital camera, respectively.

Sensing of Hg^{2+}

For selectivity experiments, 10 μL of one of the different metal ions and 10 μL of melamine in aqueous solution were mixed thoroughly first and then added into 1 mL of Au NPs solution,

the final effective concentrations of the different metal ions were all 5×10^{-6} M, whereas the final concentrations of melamine were all 1.5×10^{-6} M. After mixing thoroughly for 2 minutes, absorbance spectra and photos of Au NPs solutions at the same time were recorded with UV-Vis spectrometer and the digital camera, respectively. To study the limit of detection of Hg^{2+} , aggregation degree of 1 mL of Au NPs solution induced by melamine at a final concentration of 6×10^{-7} M was used as a control sample in the presence of 15 mM NaCl. Then 10 μL of Hg^{2+} at different concentrations and 10 μL of melamine in aqueous solution were mixed thoroughly first and then added into 1 mL of Au NPs solutions, the final effective concentrations of Hg^{2+} were ranging from 0 to 400 nM, whereas the final concentrations of melamine were all 6×10^{-7} M. After mixing thoroughly for 15 minutes, absorbance spectra and photos of Au NPs solutions at the same time were recorded with UV-vis spectrometer and the digital camera, respectively.

Results and discussion

Design of the three-input Au NPs colorimetric logic gate of Hg^{2+} , melamine, and cysteine

Fig. 1 illustrates the mechanism of the colorimetric logic gate. We define the inputs “1” and “0” of the logic gate as the presence and absence of an analyte, respectively. Then we can define the wine red (dispersed) and dark blue (aggregated) colour of Au NPs aqueous solution as “0” and “1” which correspond to the outputs of a logic gate, respectively. Since there are three kinds of analytes (Hg^{2+} , melamine, and cysteine) in our system, the possible number of input conditions is $2^3 = 8$, namely, 000, 100, 001, 101, 111, 110, 011, and 010. 13 nm colloidal Au NPs are wine-red and well dispersed in aqueous solution with the weakly surface-bound negative-charged citrate ions (input 000, output 0), whereas their aggregates appear dark blue colour. Because melamine is positively charged owing to its protonated amine groups, the addition of melamine to Au NPs solution will shield the electrostatic

repulsion of negative charges that help stabilize the Au NPs. As a result, the surface-bound citrate ions are depleted, subsequently leading to the cross link of Au NPs or the aggregation of Au NPs accompanied by a colour change from wine-red to dark blue (input 001, output 1). However, the aggregation of Au NPs induced by the addition of melamine can be easily inhibited in the presence of Hg^{2+} due to the higher affinity of melamine to Hg^{2+} , which can be explained by coordination occurred through the amine and triazine nitrogen atoms in melamine with Hg^{2+} (input 101, output 0).⁴² Among various kinds of organic groups, sulfhydryl group has the superior affinity for the surface of Au NPs due to the formation of the Au-S bond which is commonly employed in the surface modification of Au NPs.^{21, 43} As a sulfhydryl-containing amino acid, cysteine could be chemisorbed on the surface of Au NPs to form a molecular layer of cysteine (input 010, output 0). It is anticipated that the chemisorption of cysteine on Au surface is superior to the electrostatic coordination of melamine, therefore an addition of cysteine could prevent melamine from inducing the aggregation of Au NPs (input 011, output 0). On the other hand, since the breakage of Au-S bond will occur when there is Hg^{2+} in the system, the addition of Hg^{2+} to Au NPs solution in the presence of cysteine will lead to the removal of the chemisorbed thiolates on the Au surface.²⁶ This process destabilizes the well-dispersed Au NPs solution, resulting in the aggregation of Au NPs (input 110, output 1). However, the addition of Hg^{2+} to the Au NPs colloidal solution in the absence of cysteine almost has no effect on the dispersion state of Au NPs (input 100, output 0). Furthermore, in the presence of all inputs (Hg^{2+} , melamine, and cysteine), Hg^{2+} would rather bind to cysteine than melamine, this competition leads to the formation of a more stable Hg^{2+} -cysteine complex instead of the Hg^{2+} -melamine complex and subsequently resulting in the aggregation of Au NPs (input 111, output 1). Based on the coordination and ligand replacement mechanisms between Hg^{2+} , melamine, cysteine and Au NPs, the three-input colorimetric logic gate was rationally designed.




Fig. 1 A schematic illustration of the mechanism of the colorimetric logic gate.

Combinational operations of the Au NPs colorimetric logic gate

The synthesized Au NPs were characterized by TEM, indicating their size is ~13 nm (Fig. S1, ESI†). Fig. 2A shows the responses of absorption spectra of the 13 nm Au NPs solutions upon the combinational additions of melamine, cysteine, and Hg^{2+} . Dramatic changes in absorption spectra of Au NPs are observed from samples upon the additions of melamine (input 001), melamine/ Hg^{2+} /cysteine (input 111), and Hg^{2+} /cysteine (input 110). Absorption band of Au NPs in the range of 600 to ~700 nm resulted from their aggregates increases obviously while the original absorbance of Au NPs at 520 nm decreases strongly. It is found that there are insignificant changes in the absorption spectra of Au NPs upon the additions of melamine, cysteine, and Hg^{2+} in the other combinations which correspond to the input conditions 000, 100, 101, 011, and 010. TEM images of Au NPs upon the combinational additions of melamine, cysteine, and Hg^{2+} for the logic operations confirm their corresponding dispersion or aggregation states (Fig. S1, ESI†). It is consistent with the absorption spectra that the aggregates were only formed with the input conditions 001, 110, and 111. Furthermore, the corresponding colorimetric effects are evaluated by comparing the $A_{650\text{nm}}/A_{520\text{nm}}$ values (Fig. 2B). Fig. 2C depicts the network map of the combination of “AND”, “INH”, and “OR” gates for the logic operations. Neither Hg^{2+} nor cysteine can induce the aggregation of Au NPs alone, however, their coexistence would result in a severe aggregation of Au NPs. Thus, an “AND” gate could be built to convey the interrelationship between the presence of cysteine/ Hg^{2+} and the aggregation of Au NPs. Since

melamine alone also possesses the ability to enable Au NPs aggregation, an “OR” gate can be employed to describe the logic connection between melamine and the “AND” gate of cysteine/ Hg^{2+} . Meanwhile, the only presence of cysteine or Hg^{2+} can inhibit the aggregation of Au NPs induced by melamine. Therefore, two “INH” gates of melamine/cysteine and melamine/ Hg^{2+} are further needed. The detailed logic operations of “AND”, “OR”, and “INH” gates are illustrated in Table S1. Fig. 2D is a table of inputs and outputs of the combinational logic gate, the visual colour changes of the output “0” (red) and output “1” (blue) can be clearly identified. The mechanisms of analyte induced aggregation or anti-aggregation of Au NPs can be used for the discrimination of different analytes or their combinations under certain conditions. A schematic illustration of this application can be found in Fig. S2 (ESI†).

A competing coordination interaction based mechanism between Hg^{2+} , melamine and cysteine

To better understand the interactions between Hg^{2+} , melamine and cysteine, the coordination reaction between Hg^{2+} and melamine was first investigated through the titration experiments. Fig. 3A shows the absorption spectra changes of the titration of melamine solution with Hg^{2+} . The free melamine in aqueous solution exhibits a strong absorption band at 204 nm and a weak absorption band at 234 nm.^{42,44} However, upon the addition of Hg^{2+} to melamine aqueous solution, the intensity of the absorption band at 234 nm is greatly enhanced, whereas the intensity of the absorption band at 204 nm shows a pronounced




Fig. 2 Combinational logic gate operations (AND, INH, and OR) using colloidal Au NPs solution. (A) UV-vis absorption spectra responses of Au NPs solutions upon the combinational additions of melamine (6×10^{-5} M), cysteine (3.6×10^{-4} M), and Hg^{2+} (1.8×10^{-4} M). (B) The corresponding signal readouts in a function of $A_{650\text{nm}}/A_{520\text{nm}}$. (C) Network map schematically represents the combination of AND, INH, and OR gates for the logic operations. (D) The inputs and outputs of the colorimetric logic gates. Inset is the corresponding output colour of the colloidal Au NPs solutions.

decrease, indicating the coordination occurred between Hg^{2+} and melamine. The enhancement of the absorption band at 234 nm and the decrease of the absorption band at 204 nm are both observed by increasing the molar ratio of Hg^{2+} to melamine from 3:1 to 4:1, implying there is still an excess amount of nitrogen atoms which are not completely coordinated in the solution of the Hg^{2+} -melamine complex at the molar ratio of 3:1. Then a solution of the Hg^{2+} -melamine complex at the molar ratio of 3:1 was chosen to study the ligand replacement reaction of Hg^{2+} upon the addition of cysteine. Fig. 3B shows the evolution of absorption spectra of the Hg^{2+} -melamine complex at the molar ratio of 3:1 in solution upon the addition of cysteine. The absorption bands in the range of 200 to ~230 nm increase strongly and reach their maxima at the molar ratio of 1:3:6 (Melamine/ Hg^{2+} /Cysteine). To make sure this increment is not the result of the addition of cysteine itself, the equivalent amounts of cysteine were investigated solely (Fig. 3C). Compare to the increment of the band at 200 to ~230 nm induced by the addition of cysteine to the solution of the Hg^{2+} -melamine complex, the increment solely caused by cysteine is about 8 times lower, suggesting that the Hg^{2+} -melamine

complex is transformed into a more stable Hg^{2+} -cysteine complex. We also checked the absorption spectra changes of the titration of Hg^{2+} solution with cysteine at the same molar ratio ranges from 3:0 to 3:8, a similar increment of the band at 200 to ~230 nm is observed, indicating that the coordination reaction between Hg^{2+} and cysteine indeed occurred (Fig. 3D). Both of the increments of the band at 200 to ~230 nm reach their maxima when the molar ratio of Hg^{2+} to cysteine is 3:6 in the solution of Hg^{2+} -melamine and Hg^{2+} -melamine-cysteine, implying that a stable complex of $\text{Hg}^{2+}(\text{cysteine})_2$ was formed in both systems with and without melamine, which strongly suggest the complete transition of Hg^{2+} -melamine complex to Hg^{2+} -cysteine complex. It is also consistent with many previous reports that cysteine is capable of forming a complex of $\text{Hg}^{2+}(\text{cysteine})_2$ with Hg^{2+} and the stability constant of the $\text{Hg}^{2+}(\text{cysteine})_2$ complex is as high as 2×10^{40} .^{45,46} A series of control experiments were also carefully conducted and these results are evident in our proposed mechanisms that cysteine with sulfhydryl group can protect Au NPs from aggregation in the presence of melamine and can form a more stable complex with Hg^{2+} through the coordination to destabilize Au NPs




Fig. 3 Interactions between melamine, cysteine, and Hg^{2+} in aqueous solution. UV-vis absorption spectra responses of (A) melamine (2×10^{-5} M) aqueous solution upon the addition of Hg^{2+} at the molar ratio (Melamine/ Hg^{2+}) ranging from 1:0 to 1:4, (B) melamine (2×10^{-5} M) - Hg^{2+} (6×10^{-5} M) complex aqueous solution upon the addition of cysteine at the molar ratio (Melamine/ Hg^{2+} /Cysteine) ranging from 1:3:0 to 1:3:8, (C) cysteine aqueous solution at the concentration ranging from 4×10^{-5} M to 1.6×10^{-4} M, and (D) Hg^{2+} (6×10^{-5} M) aqueous solution upon the addition of cysteine at the molar ratio (Hg^{2+} /Cysteine) ranging from 3:0 to 3:8.

colloidal solution in the presence or absence of melamine (Fig S3 and S4, ESI†).

Detection of melamine, Hg^{2+} , and cysteine in aqueous solution

Au NPs colloidal solution was reported to show intrinsic colorimetric sensitivity and selectivity of to melamine,^{23, 41} we also investigated it in detail (Fig. S5, S6 and S7, ESI†). Cysteine has been widely used for inducing the aggregation of Au NPs at high concentration or low pH value for various applications due to the formation of zwitterionic networks involving head-to-head interaction of the deprotonated carboxylate ($-\text{COO}^-$) and protonated amine ($-\text{NH}_3^+$) groups on

adjacent particles.⁴⁷ In our work, we found that cysteine could also be employed as a stabilizing agent to prevent the aggregation of Au NPs induced by melamine under certain conditions. Therefore, an unusual application in which the “INH” gate of melamine/cysteine was employed in the selective and sensitive detection of cysteine was devised. Fig. 4A shows the changes in the absorption spectra of Au NPs as a result of the addition of different amino acids in the presence of melamine. It exhibits a high selectivity for cysteine over other amino acids including glycine (Gly), alanine (Ala), serine (Ser), proline (Pro), aspartic acid (Asp), asparagine (Asn), glutamic acid (Glu), tyrosine (Tyr), and arginine (Arg).




Fig. 4 Visual detection of cysteine. (A) UV-vis absorption spectra responses of Au NPs solutions upon the addition of 6×10^{-5} M different amino acids in the presence of 6×10^{-5} M melamine, (B) the corresponding signal readouts in the form of $A_{650\text{nm}}/A_{520\text{nm}}$, (C) the colour readouts of the Au NPs solutions, and (D) molecular structures of the amino acids. (E) Signal readouts in the form of $A_{650\text{nm}}/A_{520\text{nm}}$ of the Au NPs solutions upon the addition of cysteine with different concentrations in the presence of 1.4×10^{-6} M melamine, (F) the corresponding visual colour changes of the Au NPs solutions, and (G) a linear plot of $A_{650\text{nm}}/A_{520\text{nm}}$ as a function of cysteine concentration over the range of 0.6 to 1.4 μM . All the UV-vis absorption spectra and the photos of Au NPs solutions were collected after the same time (2 minutes).

acid (Glu), ornithine (Orn), glutamine (Gln), histidine (His), tyrosine (Tyr), and arginine (Arg) based on the $A_{650\text{nm}}/A_{520\text{nm}}$ values and their corresponding colour readouts (Fig. 4B and C). It can be seen from the molecular structures that only cysteine which is a sulfhydryl-containing amino acid could inhibit the aggregation of Au NPs induced by melamine (Fig. 4D). To evaluate the sensitivity of the assay, the $A_{650\text{nm}}/A_{520\text{nm}}$ values of Au NPs were measured after the addition of various concentrations of cysteine in the presence of melamine. As can be seen from Fig. 4E, the intensity ratios of the two absorption bands at 520 and 650 nm gradually decrease with the increasing amount of cysteine. Even cysteine is at the concentration of 0.6 μM , the decrease of the $A_{650\text{nm}}/A_{520\text{nm}}$ value can be still clearly obtained with the UV-vis spectrometer, demonstrating the sensitive response to cysteine. The decrease of the $A_{650\text{nm}}/A_{520\text{nm}}$ values display clear colour changes of Au NPs solutions from dark blue to wine red (Fig. 4F). As can be seen from Fig. 4G, at the cysteine concentration range of 0.6 to 1.4 μM , the $A_{650\text{nm}}/A_{520\text{nm}}$ value is closely related to the amount of cysteine added to the Au NPs solution, which can be used for the quantification of cysteine. Based on a similar strategy, the “INH” gate of melamine/ Hg^{2+} can be applied to the selective and sensitive detection of Hg^{2+} . The limit of detection for Hg^{2+} can reach as low as 50 nM (10 ppb) based on the $A_{650\text{nm}}/A_{520\text{nm}}$ value (Fig. S8, ESI†).

Conclusion

In summary, a three-input colorimetric logic gate has been designed based on the highly specific coordination and ligand replacement reactions between melamine, cysteine, Hg^{2+} , and Au NPs, and its utility has been established for visual discrimination of melamine, cysteine, and Hg^{2+} in aqueous solution. The “INH” gates of melamine/cysteine and melamine/ Hg^{2+} can also be used to quantitate cysteine and Hg^{2+} in solutions on the basis of the measurement of $A_{650\text{nm}}/A_{520\text{nm}}$ ratios of Au NPs. The “INH” gate of melamine/cysteine allows the rapid detection of cysteine down to 0.6 μM within 2 minutes. The “INH” gate of melamine/ Hg^{2+} provides a rapid detection of Hg^{2+} as low as 50 nM (10 ppb) within 15 minutes. This concept reported herein could be extended to the visual detection of a wide range of organic and biological molecules through properly designing different coordination and ligand replacement reactions based on Au NPs.

Acknowledgement

Q.X. gratefully acknowledges the financial support from Singapore National Research Foundation through its Fellowship grant (NRF-RF2009-06), and Singapore Ministry of Education via its two AcRF Tier2 grants (MOE2011-T2-2-051 and MOE2011-T2-2-085). C.C. acknowledges the support from the Central Research Support Funds of Queen's University Belfast via a start-up grant. This research is also supported in part by the CAS/SAFEA International Partnership Program for Creative Research Teams.

References

- 1 http://en.wikipedia.org/wiki/2008_Chinese_milk_scandal
- 2 <http://www.themalaymailonline.com/world/article/china-seizes-melamine-tainted-yoghurt-candy-in-new-food-scandal#sthash.quKoBjJK.dpuf>
- 3 P. M. Bolger and B. A. Schwetz, *New Engl. J. Med.*, 2002, **347**, 1735-1736.
- 4 R. Janaky, V. Varga, A. Hermann, P. Saransaari and S. S. Oja, *Neurochem. Res.*, 2000, **25**, 1397-1405.
- 5 M. PukaSundvall, P. Eriksson, M. Nilsson, M. Sandberg and A. Lehmann, *Brain Res.*, 1995, **705**, 65-70.
- 6 M. T. Goodman, K. McDuffie, B. Hernandez, L. R. Wilkens and J. Selhub, *Cancer*, 2000, **89**, 376-382.
- 7 J. K. Liu, H. C. Yeo, E. Overvik-Douki, T. Hagen, S. J. Doniger, D. W. Chu, *J. Appl. Physiol.*, 2000, **89**, 21-28.
- 8 N. Saravanan, D. Senthil and P. Varalakshmi, *Brit. J. Urol.*, 1996, **78**, 22-24.
- 9 W. Droge and E. Holm, *Faseb. J.*, 1997, **11**, 1077-1089.
- 10 I. Perlman, N. Stillman and I. L. Chaikoff, *J. Biol. Chem.*, 1940, **133**, 651-659.
- 11 W. L. Clevenger, B. W. Smith and J. D. Winefordner, *Crit. Rev. Anal. Chem.*, 1997, **27**, 1-26.
- 12 R. Kunkel and S. E. Manahan, *Anal. Chem.*, 1973, **45**, 1465-1468.
- 13 N. H. Bings, A. Bogaerts and J. A. C. Broekaert, *Anal. Chem.*, 2006, **78**, 3917-3945.
- 14 A. Bernaus, X. Gaona, J. M. Esbri, P. Higuera, G. Falkenberg and M. Valiente, *Environ. Sci. Technol.*, 2006, **40**, 4090-4095.
- 15 S. Ehling, S. Tefera and I. P. Ho, *Food Addit. Contam.*, 2007, **24**, 1319-1325.
- 16 G. M. Huang, O. Y. Zheng and R. G. Cooks, *Chem. Commun.*, 2009, 556-558.
- 17 S. Shahrokhian, *Anal. Chem.*, 2001, **73**, 5972-5978.
- 18 G. Hignett, S. Threlfell, A. J. Wain, N. S. Lawrence, S. J. Wilkins, J. Davis, R. G. Compton and M. F. Cardoso, *Analyst*, 2001, **126**, 353-357.
- 19 Y. V. Tcherkas and A. D. Denisenko, *J. Chromatogr. A*, 2001, **913**, 309-313.
- 20 C. M. Pfeiffer, D. L. Huff and E. W. Gunter, *Clin. Chem.*, 1999, **45**, 290-292.
- 21 K. L. Ai, Y. L. Liu and L. H. Lu, *J. Am. Chem. Soc.*, 2009, **131**, 9496-9497.
- 22 F. Chai, C. G. Wang, T. T. Wang, Z. F. Ma and Z. M. Su, *Nanotechnology*, 2010, **21**.
- 23 H. Chi, B. H. Liu, G. J. Guan, Z. P. Zhang and M. Y. Han, *Analyst*, 2010, **135**, 1070-1075.
- 24 J. J. Du, S. Y. Yin, L. Jiang, B. Ma and X. D. Chen, *Chem. Commun.*, 2013, **49**, 4196-4198.
- 25 J. S. Lee, P. A. Ulmann, M. S. Han and C. A. Mirkin, *Nano Lett.*, 2008, **8**, 529-533.
- 26 D. B. Liu, W. S. Qu, W. W. Chen, W. Zhang, Z. Wang and X. Y. Jiang, *Anal. Chem.*, 2010, **82**, 9606-9610.
- 27 Y. Ma and L. Y. L. Yung, *Anal. Chem.*, 2014, **86**, 2429-2435.
- 28 K. H. Su, Q. H. Wei, X. Zhang, J. J. Mock, D. R. Smith and S. Schultz, *Nano Lett.*, 2003, **3**, 1087-1090.
- 29 S. Srivastava, B. L. Frankamp and V. M. Rotello, *Chem. Mater.*, 2005, **17**, 487-490.
- 30 A. P. de Silva and S. Uchiyama, *Nat. Nanotechnol.*, 2007, **2**, 399-410.
- 31 Q. S. Mei, C. L. Jiang, G. J. Guan, K. Zhang, B. H. Liu, R. Y. Liu and Z. P. Zhang, *Chem. Commun.*, 2012, **48**, 7468-7470.
- 32 R. Freeman, T. Finner and I. Willner, *Angew. Chem. Int. Edit.*, 2009, **48**, 7818-7821.
- 33 K. S. Hettie, J. L. Klockow and T. E. Glass, *J. Am. Chem. Soc.*, 2014, **136**, 4877-4880.

ARTICLE

Journal Name

- 34 X. Q. Liu, R. Aizen, R. Freeman, O. Yehezkeli and I. Willner, *ACS Nano*, 2012, **6**, 3553-3563.
- 35 M. Zhang and B. C. Ye, *Chem. Commun.*, 2012, **48**, 3647-3649.
- 36 Z. Z. Huang, H. N. Wang and W. S. Yang, *Nanoscale*, 2014, **6**, 8300-8305.
- 37 A. Ogawa and M. Maeda, *Chem. Commun.*, 2009, 4666-4668.
- 38 Xu, X. W.; Zhang, J.; Yang, F.; Yang, X. R. Colorimetric Logic Gates for Small Molecules Using Split/Integrated Aptamers and Unmodified Gold Nanoparticles. *Chem. Commun.* 2011, **47**, 9435-9437.
- 39 D. B. Liu, W. W. Chen, K. Sun, K. Deng, W. Zhang, Z. Wang and X. Y. Jiang, *Angew. Chem. Int. Edit.*, 2011, **50**, 4103-4107.
- 40 A. Ogawa and Y. Susaki, *Org. Biomol. Chem.*, 2013, **11**, 3272-3276.
- 41 F. Wei, R. Lam, S. Cheng, S. Lu, D. A. Ho and N. Li, *Appl. Phys. Lett.*, 2010, 96.
- 42 L. L. Li, G. H. Wu, T. Hong, Z. Y. Yin and D. Sun, *Acs Appl. Mater. Inter.*, 2014, **6**, 2865-2871.
- 43 S. S. R. Dasary, A. K. Singh, D. Senapati, H. T. Yu and P. C. Ray, *J. Am. Chem. Soc.*, 2009, **131**, 13806-13812.
- 44 M. J. S. Dewar and L. Paolo, *Trans. Faraday Soc.*, 1957, **53**, 261-271.
- 45 B. V. Cheesman, A. P. Arnold and D. L. Rabenstein, *J. Am. Chem. Soc.*, 1988, **110**, 6359-6364.
- 46 W. E. Van Der Linden and C. Beers, *Anal. Chim. Acta.*, 1973, **68**, 143-154.
- 47 A. Mocanu, I. Cernica, G. Tomoaia, L. D. Bobos and O. Horovitz, *Colloids and Surfaces A: Physicochem. Eng. Aspects*, 2009, **338**, 93-101.